

ZUROLOGY

Michael P. Zahalsky, MD - Mini Varghese, MD – Christopher P. Hollowell, MD

Melissa N. Marchand, PA-C - Monika A. Kulik, PA-C – Jeffrey Bien, PA-C

5850 Coral Ridge Drive, Suite 106, Coral Springs, FL 33076

2951 N.W. 49th Avenue, Suite 308, Ft. Lauderdale, FL 33313

(954) 714-8200 Phone (954) 840-2626 fax

www.ZUROLOGY.COM



VASECTOMY

Possible Side Effects:

- Swelling of Scrotum
- Bruising of Scrotum
- Bleeding or blood clot inside scrotum
- Blood in semen
- Infection at surgery site

Complications can include:

- Fluid buildup in testicle, causing dull ache that can get worse with ejaculation
- Immune response to leaking sperm causing inflammation around testicle (sperm granuloma)
- Pregnancy in rare event (semen analysis 3 months after vasectomy will verify that no sperm is present)

Before Surgery

- Stop aspirin or any other blood thinners one week before surgery
- Shower and wash genital area thoroughly. Shave the area of surgery area
- Take medication (given to you by the office) to relax you 30 minutes to 1 hour before surgery

What to expect

- Exam of scrotum
- Numbing of surgical area
- Small incision in upper part of scrotum after being numb
- Withdraw the Vas Deferens, tie them seal them with heat and clip them
- Time of procedure is approximately 20-30minuts.

After Surgery

- Support scrotum with tight fitting underwear for at least 48 hours after surgery
- Use ice periodically for swelling and pain for the three days after surgery
- Rest for 24 hours after surgery
- Light activity for 2-3 days after surgery. Avoid sports or any lifting for approximately 1 week
- No aspirin for 3-7 days after surgery
- Avoid sexual activity for 1 week. If you do ejaculate you may notice blood in semen 3-4 days after surgery
- **Contact the office right away if you have signs of infection such as;**
 - **Pus or blood oozing from surgery site**
 - **Fever or Worsening of pain or swelling**